


African Journal of Tropical Medicine and Biomedical Research (AJTMBR)



The Journal is the Official Publication of the College of Health Sciences,
Delta State University, Abraka, Nigeria.

African Journal of Tropical Medicine and Biomedical Research (AJTMBR)
by College of Health Sciences, Delta State University is licensed
under Creative Commons Attribution-Share
Alike 4.0 International 

Editorial Board

Editor-in-Chief

Prof. Igbigbi, P. S.

Editor

Prof. Omo-Aghoja, L. O.

Associate Editors

Prof Akhator, A.

Prof Odokuma, E. I.

Desk/Managing Editor

Dr. Umukoro, E. K.

Dr. Moke, E. G.

Editorial Advisory Board

Prof Aloamaka, C. P.

Prof Asagba, S. O.

Prof. Dosumu, E. A.

Prof. Ebeigbe, P. N.

Prof Ekele, B. A.

Prof Fasuba, O. B.

Prof Feyi-Waboso, P.

Prof Ikomi, R. B.

Prof Obuekwe, O. N.

Prof Obaju-Obodo, J.

Prof Okobia, M. N.

Prof. Okonofua, F. E.

ISSN: 2141-6397

Vol. 8, No. 2, December 2025



Focus and Scope

The African Journal of Tropical Medicine and Biomedical Research is a multidisciplinary and international journal published by the College of Health Sciences, Delta State University of Abraka, Nigeria. It provides a forum for Authors working in Africa to share their research findings on all aspects of Tropical Medicine and Biomedical Sciences and to disseminate innovative, relevant and useful information on tropical medicine and biomedical sciences throughout the continent. The journal will publish original research articles, reviews, editorials, commentaries, short reports, case reports and letters to the editor. Articles are welcome in all branches of medicine and dentistry including basic sciences (Anatomy, Biochemistry, Physiology, Pharmacology, Psychology, Nursing etc) and clinical sciences (Internal Medicine, Surgery, Obstetrics and Gynaecology, Dental surgery, Child Health, Laboratory Sciences, Radiology, Community Medicine, etc). Articles are also welcome from social science researchers that document the intermediating and background social factors influencing health in countries of Africa. Priority will be given to publication of articles that describe the application of the principles of primary health care in the prevention and treatment of diseases.

Editorial Notices

The journal will be published biannually in the months of March and September. Annual subscription fee in Nigeria is two thousand naira (N2,000) per volume (2issues); One-thousand-naira single copy (N1000). The annual subscription rate for other parts of the world is as follows: United Kingdom £60 (post free). West Africa \$60 (post free). The rest of the World and the United States of America \$120 (post free). A charge of \$60 is made for reprints inclusive of postage. Cheques should be made payable to the African Journal of Tropical Medicine and

Biomedical Research and addressed to the Editor-in-Chief.

Journal Contact

All correspondence, including manuscripts for publication (in triplicate) should be addressed to:

Professor P.S. Igbigbi

The Editor-in-Chief,
Department of Anatomy,
Faculty of Basic Medical Sciences,
College of Health Sciences,
Delta State University, Abraka,
Delta State, Nigeria.

Or:

Professor Lawrence Omo-Aghoja

Editor
Department of Obstetrics and
Gynecology,
Faculty of Clinical Medicine,
Delta State University, Abraka, Nigeria.
Email: journalajtmbr@yahoo.com
Cc: all email to
eguono_2000@yahoo.com
Tel: 08039377043

All authors are advised to submit an electronic copy in CD-ROM along with a hard copy of their manuscript, as this will spare remarkable time in the reviewing and typesetting processes.

In the alternative, authors can submit their articles and covering letter by email attachments. A covering letter (signed by all authors) accompanying the manuscript should certify that the article has not been previously published and is not being considered for publication elsewhere.

Information for Authors

All manuscript are peer-reviewed and accepted with the understanding that the work has not been published or being considered for publication elsewhere. Indeed, the authors would be requested

to sign a copyright form transferring the ownership of the paper to the African Journal of Tropical Medicine and Biomedical Research. All articles must include the correct names and addresses of author(s) including e-mail addresses and telephone numbers. Articles will be subjected to a thorough peer review process before any decision is made to publish or not. Authors should note that the African Journal of Tropical Medicine and Biomedical Research is not under any obligation to publish articles submitted, as decision to publish will be based on recommendations of reviewers and the editorial advisory board.

Manuscripts

Articles submitted for publication should be typed double-spaced with 2.5cm margins with accompanying CD-ROM in Microsoft Word format for easy and quick peer review and typesetting. Each of the following sections should begin in a new page: title page, abstract, introduction, materials and methods, results, discussion, acknowledgment (s), references, tables, legends to figures and illustrations. The manuscript should include:

Title Page

The title page should include the following information: 1. the title and sub-title; 2. the name(s) of the author(s); 3. the affiliation(s) of the author(s); 4. name and address of the corresponding author and 5. three to six key words for indexing and retrieval purposes.

Abstract

The abstract should be structured and not more than 250 words. It should carry the following headings: Introduction, Materials and Methods, Results and Conclusion.

Original Research- The journal welcomes

articles reporting on original research, including both quantitative and qualitative studies. Full-length articles should generally not exceed 3000 words, excluding abstract, tables, figures, and references. The subject matter should be organised under appropriate headings and sub-headings as itemized above.

Review Articles- Comprehensive review articles on all aspects of tropical medicine and biomedical sciences will also be considered for publication in the journal. Reviews should provide a thorough overview of the topic and should incorporate the most current research. The length of review articles must not exceed 3,000 words and the organisational headings and sub-headings used are at the author's discretion.

Short Reports - Brief descriptions of preliminary research findings or interesting case studies will be considered for publication as short reports. The length of the abstract and article should be restricted to 150 and 2,000 words respectively and organisation of short reports are left to the author's discretion.

Commentaries or Editorials- Commentaries or editorials on any aspect of tropical medicine and biomedical sciences in Africa will be considered for publication in the journal. Opinion pieces need not reference previous research, but rather reflect the opinions of the author(s). The length should not exceed 2,000 words.

Tables and Figures

All tables and figures should be submitted on separate sheets of paper and should be clearly labelled. Coloured tables and figures may be reprinted in black and white. Authors should especially take care that all tables are clear and understandable by themselves, independent of

the text. A reader should be able to read only the tables and easily grasp all information without the text.

Acknowledgments

Acknowledgments should be included on a separate sheet of paper and should not exceed 100 words. Funding sources should be noted here.

References

References should be in the Vancouver style and numbered consecutively in the order in which they are mentioned in the text. Titles of journals should be abbreviated according to the Index Medicus style. Authors must cross-check and make sure that all information provided in the reference list is complete and correctly written. Reference numbers should be inserted above the line on each occasion a reference is cited in the text, e.g., ... as 1-3 reported in other studies. Numbered references should appear at the end of the article and should include the names and initials of all authors. The format of references should be as published by the International Committee of Medical Journal Editors in the British Medical Journal 1988, volume 296, pages 401-405. The following are sample references

for an article published in a journal and for a book: Ahmed Y, Mwaba P, Chintu C, Grange JM, Ustianowski A, Zumla A. A study of maternal mortality at the University Teaching Hospital, Lusaka, Zambia: the emergence of tuberculosis as a major non-obstetric cause of maternal death. *Int J Tuberc Lung Dis* 1999; 3: 675-680. Whitby LG, Smith AF, Beckett GJ. *Enzyme Tests in Diagnosis*. In: *Lecture Notes on Clinical Chemistry*. Whitby LG, Smith AF & Beckett GJth (eds). 4 editions. Blackwell Scientific Publications. 1988. 103-127.

Units of Measurement

All measurements should be expressed in SI (Systeme International) Units.

Galley proofs

Corrections of galley proofs should be strictly restricted to Printer's error only. Orders for offprints should be made when the corrected proofs are being returned by the authors. Articles accepted for publication remain the property of the journal and can only be reproduced elsewhere in line with section 5 of the copyright agreement.

“For those journals with their own websites, and a very high degree of in-house technical competency, it is possible to get code snippets from the Creative Commons website for the license type chosen, which can be embedded in your online content, but please note that that is definitely not a requirement, just a “nice to have”.




Table of Contents

Editorial

- The Role of Forensic Medicine in Criminal Justice Delivery in Nigeria 7-11
Umeaku, U

Original articles

- Differential Alterations In Behavioural Phenotypes, Brain Biochemical Profiles And Histomorphology In Mice After Administration Of Sildenafil-dapoxetine Formulation 12-26
Omogbiya A, Orowigbo O, Moke EG, Obob M, Emudainobwo JOT, Awbin PE, Saviour GU
- Pre-Examination Stress Among Pre-Clinical Medical Students: A Nigerian Survey 27-33
Ominde, BS, Enaobwo, MT, Jeremiah, O, Omoro, OF, Igbigbi, PS.
- Evaluation Of Computational And Insecticidal Activities Of Oils From *Ocimum Gratissimum* And *Cymbopogon Citratus* Against *Anopheles Gambiae* Mosquito 34-49
Elelegu, EJ, Dunkwu, CC, Enyi, KC, Onuelu, JE, Onyesom, I.
- Induced Abortion As A Method Of Contraception By Married Women With Unintended Pregnancy In Delta State, Nigeria: Prevalence, Predictors, And Reproductive Health Burden. 50-60
Williams, OO, Ayo, N, Omonigbo, E.
- Duration-Dependent Post-Trauma Sleep Deprivation Differentially Modulates PTSD-Like Anxiety and Oxidative Markers in Mice 61-70
Ben-Azu B, Omogbiya AI, Ayereoghene SM, Usin SG, Oyonmi MO
- Assessment of cytotoxicity and growth inhibitory effects of methanol extract of *Ageratum conyzoides* Linn 71-77
Ikoya, S, Apitikori-Owumi, JE, Nwoguezè, BC, Agboola, OE, Ekakitie, LI, Odegbe, OJ, Ofoke, IH, Oviri, MO.
- Evaluating The Impact Of Computer Assisted Cognitive Remediation On Recovery Outcomes In Schizophrenia: A Quasi Experimental Study From Northwest Nigeria. 78-85
Yakubu AA, Bashir SM, Aweka AI, Murinyi MM, Khalid S, Yakasai BA
- Biochemical Impact Of Occupational Cement Dust Exposure On Block Industry Workers In Rivers State, Nigeria. 86-90
John, GC, Anosike, IK

Induced Abortion As A Method Of Contraception By Married Women With Unintended Pregnancy In Delta State, Nigeria: Prevalence, Predictors, And Reproductive Health Burden

¹Oduvbun WO, ²Nwachi A, ³Esemuede O.

ABSTRACT

Introduction: Unsafe abortion is an important cause of maternal death, globally. Nigeria contributes about twenty percent to the global figure. Induced abortion by married women as a method of contraception may be associated with life threatening complications. The study determined the prevalence, predictors, and the reproductive health burden of induced abortion, as a method of contraception, among married women with unintended pregnancy in Delta State.

Materials and Method: This was a cross-sectional study conducted in three public health facilities in Delta State. Participants were recruited through census sampling. Interviewer assisted questionnaire was used to collect data, which was subsequently analysed Statistical Product and Service Solution version 23. The level of statistical analysis was set at $P < 0.05$.

Results: The age range of participants was 18-49 years, with a mean age of 29.52 ± 5.78 years. Prevalence of unintended pregnancy was 58.0%. A third (30.7%) had a history of contraceptive use, while 42.3% a history of termination of pregnancy. The major (38.8%) reason for unintended pregnancy was contraceptive failure. The main (8.8%) complication was abdominal pain. The number of living children and unplanned pregnancy were the predictors for induced abortion ($P < 0.001$; $P < 0.001$), respectively.

Conclusion: This study showed a high prevalence of induced abortion among multiparous married women with unintended pregnancy, in Delta State. However, there was minimal adverse reproductive health outcome among the participants.

1. Department of Obstetrics and Gynaecology, College of Health Sciences, Delta State University, Abraka. <https://orcid.org/0000-2-0828-3215>

2. Department of Obstetrics and Gynaecology, Central Hospital, Warri, Delta State, Nigeria. Nwachi4aya@gmail.com

3. Department of Obstetrics and Gynaecology, Eku Baptist Government Hospital, Delta State, Nigeria. esemuedeomonigbo@yahoo.com

Corresponding author: Williams Obukohwo Oduvbun, Department of Obstetrics and Gynaecology, College of Health Sciences, Delta State University, Abraka. willymeg1@gmail.com +2348033722401.

INTRODUCTION

Induced abortion is a medical or surgical procedure that allows artificial termination of unintended or unwanted pregnancies.¹ On a global scale, 25 million of unsafe abortion occur annually, with estimated 97% taking place in developing countries.² Unsafe abortion is among the leading causes of maternal mortality and morbidity in developing countries.^{3,4}

In Nigeria, and most part of Sub-Saharan Africa, induced abortion is illegal and considered criminal offence, with resultant procurement of abortions by women in clandestine, unhygienic environments, and adverse outcome, such as haemorrhage, pelvic sepsis, genital tract trauma, uterine perforation among others.^{3,5,6} The long-term reproductive health complication include chronic pelvic inflammatory disease, ectopic

pregnancy and infertility.⁷

A clinical study by Oye-Adeniran et al,⁸ in the Western part of Nigeria, showed that 30.2% of women who presented for induced abortion were married women, and the aim of their study was to determine the use of pregnancy termination for birth spacing or limitation of births. This practice questions the adoption and practice of modern or even traditional methods of contraception in the study setting and at the time of study. Our study however, aimed to determine the prevalence, predictors, and the reproductive health burden of induced abortion by married women with unintended pregnancy in Delta State, South-South Nigeria. To the best knowledge of the authors, there are no studies on induced abortion by married women in Delta State. It is hoped that this multicentred-health-facility study, would give some insight into the behavioural pattern of married women with unintended pregnancy, contraceptive usage, and induced abortion in Delta State. Findings from this study may help in shaping policy formulation by the state Government on birth control.

MATERIALS AND METHODS

This was a cross-sectional survey conducted in the following three health facilities in Delta State: The Delta State University Teaching Hospital (DELSUTH), Oghara. This is a tertiary health facility, sited in the semi-urban community of Oghara, in the Ethiope-West Local Government Area; Central Hospital Warri (CHW), is a tertiary Health facility located in the densely populated city of Warri, which is regarded as the commercial State Capital, and located in Warri South Local Government Area; Eku Baptist Government Hospital (EBGH), is a busy rural Secondary Health facility, located in the Ethiope East Local Government Area. While DELSUTH and EBGH are located in the

Delta Central Senatorial zone, CHW is located in Delta South Senatorial zone. The study population were allocated proportionately according to the different monthly post- natal volume of married women between the age of 18 and 49 years. The CHW, had the largest population of respondents, followed by EBGH. The study was conducted in three months, from January 1, 2025 - March 31, 2025.

At each facility, eligible respondents were recruited consecutively in a census sampling technique until the allotted number for each facility was completed.

Inclusion criteria: Married women from the age of 18-49 years, with at least a living child who presented at the post- natal clinic of study facilities.

Exclusion criteria: Single mothers and acutely ill-married women who required emergency service Data was collected using a semi-structured interviewer administered questionnaire on sociodemographic characteristics, reproductive health history and knowledge. Information was obtained on unintended pregnancy and the induced abortion. The population in the study settings widely speak and understand the English Language, and where indicated interpreters were deployed. Information from the study questionnaires was transferred to a computer dataset, which was cleaned up by the lead researcher before data analysis.

Data processing and Analysis.

Data collected was analysed using SPSS version 25.0 software. The socio-demographic characteristics of respondents was summarized using frequency tables and percentages. The awareness of respondents about contraceptives; the use of contraceptives; desire to space and limit birth; proportion of married women with

unintended and induced abortion was, method of induced abortion, and complication from induced abortion was determined by using univariate analysis. Chi-square was used for test of association; a p-value of <0.05 was considered statistically significant. Multivariate binary logistic regression was used to assess the predictors of induced abortion.

RESULTS

A total of 390 eligible women were recruited for the study, out of which the questionnaires for 362(92.8%) was available for analysis.

Information in the remaining 28 proformas was insufficient for inclusion in the analysis.

Table 1 shows the baseline sociodemographic parameters of the study population. Many (53.6%), of the participants were in the 21-30 age group, with a mean age of 29.52 ± 5.78 years. In addition, the majority (95.6%), of the participants were Christians, with secondary level of education (50.6%), traders (48.9%), and had 1 child (35.4%).

Table 1: Sociodemographic parameters of participants

		Frequency (n)	Percentage (%)
<i>Age group (years)</i>	≤ 20	21	5.8
	21-30	194	53.6
	31-40	138	38.1
	≥ 41	9	2.5
	<i>Mean \pm SD</i>	29.52 ± 5.78	
<i>Religion</i>	<i>Christian</i>	349	95.6
	<i>Islam</i>	12	3.3
	<i>Others</i>	4	1.1
<i>Level of Education</i>	<i>No formal education</i>	6	1.6
	<i>Primary</i>	26	7.2
	<i>Secondary</i>	183	50.6
	<i>Post-secondary</i>	147	40.6
<i>Occupation</i>	<i>Unemployed</i>	57	15.7
	<i>Unskilled</i>	32	8.8
	<i>Artisan</i>	35	9.7
	<i>Trader</i>	177	48.9
	<i>Farmer</i>	12	3.3
	<i>Professional</i>	49	13.5
<i>Number of children</i>	1	128	35.4
	2	109	30.1
	3	68	18.8
	4	41	11.3
	5	16	4.5

The prevalence of unintended pregnancy and induced abortion is shown in Table 2. Only about a third of the participants (30.7%) had ever used any contraceptive, with more than half (58.0%) having unintended pregnancies.

Furthermore, 42.3% of the participants had induced abortion, with 21.0% performing it once, while 15(6.9%) participants had a history of three or more induced abortions.

Table 2: Prevalence of unintended pregnancy and induced abortion among participants

		Frequency (n)	Percentage (%)
<i>Contraceptive ever used</i>	<i>Yes</i>	111	30.7
	<i>No</i>	251	69.3
<i>Unintended pregnancies</i>	<i>Yes</i>	210	58.0
	<i>No</i>	152	42.0
<i>Number of Unintended Pregnancies</i>	<i>None</i>	152	42.0
	<i>1</i>	68	18.8
	<i>2</i>	75	20.7
	<i>3</i>	34	9.4
	<i>≥4</i>	33	9.2
<i>History of Induced abortion</i>	<i>Yes</i>	153	42.3
	<i>No</i>	209	57.7
<i>History of Number of induced abortions (n=153; 42.3%)</i>	<i>1</i>	76	21.0
	<i>2</i>	52	14.4
	<i>≥3</i>	25	6.9

Table 3 shows the reasons for unintended pregnancy and actions taken by participants. The commonest reason for unintended pregnancy was Contraceptive failure (38.8%) and of this group of participants, Emergency contraceptives failure was the leading (25.7%) cause. The commonest method of induced abortion was the use of drugs (34.3%). Other

methods of induced abortion by participants, included MVA (8. %) and D&C (11.3%),

In addition, of the 53.6% participants who had induced abortions, 17.4% (63/362) were successful, while 36.2% (31/362) were unsuccessful in the attempts at the termination. Sixteen (4.4%), participants did nothing about the unplanned pregnancies

Table 3: Reasons for unintended pregnancy, method of contraception, actions taken by respondents and method of termination

		Frequency (n)	Percentage (%)
<i>Reasons for unintended pregnancy</i> (n=210; 58.0%)	No contraceptive use	40	19.2
	Contraceptive failure	170	38.8
<i>Methods of contraceptive that failed</i> (n=170; 47.0%)	Implant	11	3.0
	Daily pills	27	7.5
	IUCD	2	0.6
	Natural method	21	5.8
	Injectables	16	4.4
	Emergency contraceptives	93	25.7
<i>Actions taken concerning the unplanned pregnancy</i> (n=210; 58.0%)	Unsuccessful attempt at termination **	63	17.4
	Successful attempt of the unplanned pregnancy **	131	36.2
	<i>Did nothing</i>	16	4.4
<i>Methods of induced abortions</i>	<i>Drugs</i>	124	34.3
	<i>MVA</i>	29	8.0
	<i>D & C</i>	41	11.3

** Total number of induced abortions = 194 (53.6%) participants

Table 4 shows the personnel responsible for the induced abortion, and pattern of post-abortion complications. The majority 27.3% of study participants who carried out induced abortion was by Misoprostol purchased at patent medicine (Chemist) stores, and 16% of

respondents with history of induced abortion suffered post-abortion complications, with the commonest complication being abdominal pain. The complications were mostly (27/362), treated in hospitals/clinic

Table 4: Personnel responsible for termination, pattern of post-abortion complications and treatment of the complication.

		Frequency (n)	Percentage (%)
<i>Who performed the abortion</i> (n=194; 53.6%)	Doctors	9	2.5
	Nurse	34	9.4
	Pharmacist	18	5.0
	Self-administered	99	27.3
	Misoprostol from Chemist stores		
	Trad. Birth Attendants	34	9.4
<i>Place where the termination took place (n=194; 53.6%)</i>	<i>Private Health Facility</i>	47	12.9
	<i>Chemist</i>	69	19.1
	<i>Trad. Birth Attendants</i>	34	9.4
	<i>Others</i>	44	12.2
<i>Post abortion complication</i>	<i>Yes</i>	58	16.0
	<i>No</i>	136	37.6
<i>Type of Complication</i> (n=58; 16.0%)	<i>Abdominal pain</i>	32	8.8
	<i>Excessive Bleeding</i>	26	7.2
<i>Treatment of Post abortion complications (n=58; 16.0%)</i>	<i>Hospital/ clinic</i>	27	7.5
	<i>Chemist</i>	16	4.4
	<i>Self-medication</i>	15	4.1

Table 5 highlighted the association between the sociodemographic and induced abortion. A comparison between the sociodemographic parameters with induced abortion showed that there was a statistically significant association when comparing the age groups, religion, occupation and number of children with the

study outcome ($p < 0.050$). Comparing the age groups, religion, occupation and number of children between those with induced abortion and those without, revealed that more participants in the former group were between 31-40 years (42.8% vs. 32.1%), were Christians (94.8% vs. 96.4%) and traders (51.0% vs. 46.4%).

Table 5: Association between socio-demographic and induced abortion

		Induced abortion N (%)	No abortion N (%)	χ^2	p-value
<i>Age group (years)</i>	<i>≤20</i>	10 (5.2)	11 (6.5)	9.686	*0.021
	<i>21-30</i>	93 (47.9)	101 (60.1)		
	<i>31-40</i>	83 (42.8)	55 (32.7)		
	<i>≥41</i>	8 (4.1)	1 (0.6)		
<i>Religion</i>	<i>Christian</i>	184 (94.8)	162 (96.4)	8.911	*0.012
	<i>Islam</i>	10 (5.2)	2 (1.2)		
	<i>Others</i>	0 (0.0)	4 (2.4)		
<i>Level of Education</i>	<i>None</i>	3 (1.5)	3 (1.8)	0.200	0.978
	<i>Primary</i>	14 (7.2)	12 (7.1)		
	<i>Secondary</i>	100 (51.5)	83 (49.4)		
	<i>Post-secondary</i>	77 (39.7)	70 (41.7)		
<i>Occupation</i>	<i>Unemployed</i>	35 (18.0)	22 (13.1)	7.486	0.187
	<i>Unskilled</i>	13 (6.7)	19 (11.3)		
	<i>Artisan</i>	21 (10.8)	14 (8.3)		
	<i>Trader</i>	99 (51.0)	78 (46.4)		
	<i>Farmer</i>	5 (2.6)	7 (4.2)		
	<i>Professional</i>	21 (10.8)	28 (16.7)		
<i>Number of living children</i>	<i>1</i>	23 (11.9)	105 (62.5)	102.382	*<0.001
	<i>2</i>	77 (39.7)	32 (19.0)		
	<i>3</i>	53 (27.3)	15 (8.9)		
	<i>4</i>	30 (15.5)	11 (6.5)		
	<i>5</i>	11 (5.7)	5 (3.0)		

Table 6 shows that the number of living children and history of unplanned pregnancies were predictors of induced abortion by married women, $P < 0.001$ and $P < 0.005$, respectively.

Table 6: Regression analysis to show the predictors of induced abortion.

Categories		B	S.E.	OR	95% C.I. for OR		p-value
					Lower	Upper	
Age Group	≤ 20			1.000			0.690
	21-30	0.361	0.544	1.435	0.494	4.168	0.507
	31-40	0.123	0.572	1.130	0.369	3.466	0.830
	≥ 41	-0.501	1.253	0.606	0.052	7.067	0.689
Religion	Christian			1.000			0.699
	Islam	-1.105	1.260	0.471	0.032	4.307	0.999
	Others	-1.246	1.733	0.398	0.021	8.974	0.999
Number of living children		-0.729	0.129	0.482	0.375	0.621	* <0.001
History of contraceptive use		0.351	0.279	1.420	0.821	2.455	0.210
History of unplanned pregnancy		1.503	0.265	4.493	2.674	7.551	* <0.001

Discussion.

This study examined the prevalence, predictors, and reproductive health burden of induced abortion by married women with unintended pregnancy, as a method of contraception.

There was a very high (58%) rate of unintended pregnancies, and a prevalence rate of 42.3% of induced abortion by married women in the study. There was 30.7% contraceptive uptake by the participants. Contraceptive failure (38.8%) was the main reason for the unplanned pregnancies. The use of drugs was the popular (34.3%) method of termination, while abdominal pains (8.8%) was the leading complication. The number of living children and history of unplanned pregnancies were predictors of induced abortion by participants.

The high rate of unintended pregnancy in this study higher than rates in similar studies in the western part of Nigeria.⁹ The rate was also

higher than the 41.4%,¹⁰ in a study in Ethiopia. The differences may be related to the heterogeneity in contraceptive awareness and uptake in the various settings. Unplanned pregnancies are frequently associated with induced abortions.^{11,12}

Contraceptive failures were the leading cause of unplanned pregnancy in this study. The contraceptive failure rate observed in this was considerably higher than that reported in a survey conducted across 8 Nigerian states,¹³ where 16% of women using modern contraceptive methods, mainly daily oral pills, reported unintended pregnancies. In contrast, emergency contraceptive use was the predominant contributor to contraceptive failure in the present study. While, Emergency contraceptive pills may be considered a useful backup method of pregnancy prevention, its use as a primary method of birth control may result in unplanned pregnancies as shown in this study.

The prevalence rate of induced abortion in this study was considerably higher than the rate by Ojo, et al, and Okonofua, et al.^{9, 14}. In the study by Okonofua, the participants were made up of both single and married women, while Ojo, et al used the same denominator in this study, used a combination of quantitative and qualitative methods. The differences in prevalence rates may be a reflection of the study designs, settings and social behaviour spanning different timelines.

The patronage of patent medicine dealers in this study for the conduct of pregnancy termination by a high number of participants was an unusual observation. This finding contrasted with the 82% documented by Ojo, et al,⁹ in the western part of Nigeria, and 45.6%,¹⁵ in India, in which majority of induced abortions were carried out in private health facilities. The widespread distribution of chemist stores in part of Nigeria where this study was conducted and, across the counter sale of Misoprostol for medical termination of pregnancy may have been responsible for this observation, and likely explain the difference between the result of this study and that by Ojo et al.⁹ Nigeria, like many other countries in the Sub-Saharan region of Africa have restrictive abortion laws.^{3, 16} Because of the legal consequences of pregnancy termination in public health facilities in Nigeria and other parts of Sub-Saharan African countries, most women in need of abortion services go about it in clandestine manner, including patronising Chemist stores for drugs.³ The use of Misoprostol obtained from medicine stores appears to be one of the ways to circumvent the restrictive abortion laws. The complications of abdominal pains and bleeding observed among the participants who had induced abortion is similar to previous studies.^{9, 14, 17} Treatment of complications were mostly in private or public health facilities. None

of the participants suffered any serious adverse outcome from the induced abortion. While the study did not include the gestational age at which the pregnancies were terminated, the possibilities of early pregnancy in most of the participants could not be excluded.

There was a significant association between the older (≥ 31 years) married women and induced abortion. A previous study,⁹ documented that married women who are 40 years and beyond had three-fold likely to terminate their pregnancy, compared to the younger age groups. It could be argued that the need to limit family size, the need to reduce social -economic burden in raising a large family, among others, may be an explanation for induced abortions in the older women, who are also of higher parity. Association between increasing parity and induced abortion in this study was also demonstrated in other studies in Nigeria, Kenya, and in India.^{9, 18, 19, 20}

Unplanned pregnancies, especially, among women who are multiparous were predictive of induced abortion in this study. For several women with unintended pregnancies, there may be other undisclosed reasons underpinning their decisions to terminate another pregnancy that was not desired.

Conclusion

This study showed a high prevalence of induced abortion among multiparous married women with unintended pregnancy, in Delta State. However, there was minimal adverse reproductive health outcome among the participants that had induced abortion. The need for more effective education of women during prenatal and post-natal clinic on contraceptive effectiveness and provision is suggested by this study.

Study limitations

This study has some limitations, which included the small sample size; the exclusion of 28 (7.2%) questionnaires may have affected the completeness of the data that was analyzed. Considering the illegal status of abortion in Nigeria, it is impossible to exclude some level of under-disclosure of induced abortion among participants. It is hoped that the confidentiality and the anonymity assurance, would have addressed the fear of participants' prosecution. Another study limitation was the fact that the study took place in two out of the three senatorial zones in Delta State, thus making the generalisation of study findings difficult. These two zones are largely populated by the Urhobos, Itsekiris, and the Ijaw ethnic groups. Married women in the Delta North senatorial zone, inhabited by the Ibo speaking ethnic group, were not included in this study. Further study studies may be required, that is more inclusive, that should include qualitative components that may capture some socio-cultural nuances that could contribute to induced abortion by married in Delta State.

Acknowledgement

On behalf of the authors, I want to thank all the staff of the study facilities and express my appreciation to the study participants for their co-operation during the conduct of the student.

REFERENCES

1. Wei Z, Yu D, Liu H. Induced abortion among married women in China: a study based on the 1997-2017 China Fertility Surveys. *China Popul Dev Stud.*2021;5(1):83-99.doi:10.1007/s42379-021-00079-5.
2. Ganatra B, Gerdtz C, Rossier C, Johnson BR, Tuncalp O, Assfi A, et. Global, regional, and subregional classification of abortions by safety, 2010-14: estimates from a Bayesian hierarchical model. *Lancet.* 2017; 390(10110):2372-2381.doi: 10.1016/ S0140-6736(17)31794-4.
3. Odunvbun WO, Kollie JT. The burden of surgical complications from unsafe abortion treated at the John F. Kennedy Maternity Center, Monrovia. Liberia. *Malawi Med.J.* 2022;34(1):43-48.
4. Omideyi AK, Akinyemi AI, Aina OI, Adeyemi CA, Fadeyibi OA, Bamiwuye SO, et al. Contraceptive practice; unwanted pregnancies and induced abortion in Southwest Nigeria. *Global Public Health.* 2011;6(Suppl 1):S52-S72.
5. Bankole A, Adewole IF, Hussain R, Awolude O, Singh S, Akinyemi JO. The incidence of abortion in Nigeria. *Int Perspect Sex Reprod Health.* 2015;41(4):170.
6. Oye-Adeniran BA, Umoh AV, Nnatu SN. Complications of unsafe abortion: a case study and the need for abortion law reform in Nigeria. *Reprod Health Matters.* 2002;10(19):18-21.
7. Shah PS, Zao J. Induced termination of pregnancy low birthweight and preterm birth: a systematic review and meta-analyses. *BJOG.* 2009;116(1):1425-1442.
8. Oye-Adeniran BA, Adewole IF, Umoh AV, Fapohunda OR, Iwere N. Characteristics of abortion care seekers in South-Western Nigeria. *Afri J Reprod Health* 2004:81-91.
9. Ojo IE, Ojo TO, Orji EO. Why do married women procure abortion? Experiences from Ile-Ife, Southwestern Nigeria. *Afr Health Sci.* 2021; 21(1): 327-337. [Doi:10.4314/ahs.v21i1.42](https://doi.org/10.4314/ahs.v21i1.42)
10. Geda NR, Lako TK. A population-based study on Unintended pregnancy among married women in a district in Southern Ethiopia. *J Geogr Reg Plan.* 2011;4(7) :417.
11. Wang H, Zou Y, Liu H, Chen X. Analysis of unintended pregnancies and influencing factors among married women in China.

- Cbna Popul Dev Stud.* 2023;7:15-36.
12. Bearak J, Popinchalk A, Ganatra B, Moller AB, Tuncalp O, Beavin C, et al. Unintended pregnancy and abortion by income, region, and the legal status of abortion: estimates from a comprehensive model for 1990-2019. *Lancet Global Health.* 2020;8(9):e1152-e1161.
 13. Sedgh G, Bankole A, Oye-Adeniran B, Adewole IF, Singh S, Hussain R. Unwanted pregnancy and associated factors among Nigerian women. *Int Fam Plann Perspect.* 2006;175-184.
 14. Okonofua FE, Odimegwu C, Ajabor H, Daru PH, Johnson A. Assessing the prevalence and determinants of unwanted pregnancy and induced abortion in Nigeria. *Studies in Family planning.* 1999;30(1):79-83.
 15. Dhillon BS, Chandhiok N, Kambo I, Saxena NC. Induced abortion and concurrent adoption of contraception in the rural areas of India (an ICMR task force study) *Indian Med Sci.* 2004; 58:478-84.
 16. Criminal Code Act. Chapter 77 of the Laws of the Federation of Nigeria (Revised ed. 1990), Articles 228-230,297, 309,328.
 17. Lamina M. Prevalence and determinants of unintended pregnancy among women in south-western Nigeria. *Ghana Med J.* 2015;49(3):187-194.
 18. Yaya S, Amouzou A, Olalekan AU, Ekholuenetale M, Bishwajit G, Udenigwe O, et al. Prevalence and determinants of terminated and unintended pregnancies among married women: analysis of pooled cross-sectional surveys in Nigeria. *BMJ Global Health* 2018; 3(2): e000707.
 19. Maina BW, Mutua MM, Sidz EM. Factors associated with repeat induced abortion in Kenya. *BMC Public Health.* 2015; 15(1):1048
 20. Rana J, Sen KK, Sultana T, Hossain MB, Islam RM. Prevalence and determinants of menstrual regulation among ever-married women in Bangladesh: evidence from a national survey. *Reprod health.* 2019;16(1):123.

Odunvbun WO, Nwachi A, Esemuede O. . Induced Abortion As A Method Of Contraception By Married Women With Unintended Pregnancy In Delta State, Nigeria: Prevalence, Predictors, And Reproductive Health Burden. *Afr. J. Trop. Med. & Biomed. Res.* 2025; 8(2) 50-60
<https://dx.doi.org/10.4314/ajtmbr.v8i2.4>